

## ANTIDIABETIC ACTIVITY OF METHANOLIC EXTRACT OF LEAF OF *BAUHINIA PURPUREA*

S. PAHWA<sup>\*1</sup>, R. MAZUMDER<sup>1</sup>, S. BHATTACHARYA<sup>2</sup>

<sup>\*1</sup>Department of Pharmaceutical Technology, Noida Institute of Engineering and Technology, 19, Knowledge Park-II, Greater Noida-201306, Uttar Pradesh, India

<sup>2</sup> Shambhunath Institute of Pharmacy, Jhalwa, Peepal Gaon, Allahabad-211012, Uttar Pradesh, India

### ABSTRACT

Diabetes mellitus is a major disease characterized by derangement in carbohydrate, fat and protein metabolism, affecting nearly 10% of the population. In the recent past many hypoglycaemic agents have been introduced, still diabetes and the related complications continue to be a major medical problem, not only in developed countries but also in developing countries. Many Indian medicinal plants are reported to be useful in diabetes. However, search for new antidiabetic drugs continues.

*Bauhinia purpurea* (family: Caesalpiniaceae) is a medium sized, evergreen, ornamental tree found throughout India. The aerial parts of the plant are reported to contain flavanone glycosides, foliar flavonoids, 6-butyl-3-hydroxy flavanone, amino acids, phenyl fatty ester, lutine and  $\beta$ - sitosterol. Flavonoids are polyphenolic compounds that are widely distributed in the plant kingdom. They are reported to exhibit various pharmacological activities such as CNS activity, cardiotoxic activity, lipid-lowering activity, antioxidant activity, hepatoprotective activity and **hypoglycemic** activity.

The present investigation was carried out to study the antidiabetic effect of the methanolic extract of leaf of *Bauhinia purpurea* in Streptozotocin (STZ) induced diabetic model. The antidiabetic activity was evaluated in normal & STZ induced diabetic rats. Decreased blood glucose levels of the test animals showed that the extract exhibited significant antidiabetic activity, when compared to diabetic control group. The results also indicated dose dependent effect. Vehicle control animals were found to be stable in their body weight, but diabetic rats showed significant reduction in body weight during 15 days. The effect of methanolic leaf extract of *Bauhinia purpurea* on glucose level in STZ induced diabetic rats exhibited a significant reduction ( $P < 0.01$ ) in blood glucose level, when

compared to the diabetic control group. The antidiabetic activity produced by the extract might be due to the increased uptake of glucose at the tissue level or due to an increase in pancreatic beta cell function or due to inhibition of intestinal absorption of glucose.

**Keywords:** *Bauhinia purpurea*; Bark; Methanolic extract; Antidiabetic; Streptozotocin

### INTRODUCTION:

Diabetes mellitus (DM) is one of the common metabolic disorders with micro and macrovascular complications that results in significant morbidity and mortality. It is considered as one of the five leading causes of death in the world.<sup>1,2</sup> In modern medicine no satisfactory effective therapy is still available to cure diabetes mellitus.<sup>3</sup> There is increasing demand by patients to use natural products with antidiabetic activity due to side effects associated with the use of insulin and oral hypoglycemic agents. There are numerous traditional medicinal plants reported to have hypoglycemic properties such as *Allium sativum* (Garlic), *Azadirachta indica* (Neem), *Vinca rosea* (Nayantara), *Trigonella foenum* (Fenugreek), *Momordica charantia* (Bitter ground), *Ocimum santum* (Tulsi) and so on. Many of these are less effective in lowering glucose levels in severe diabetes.<sup>4-6</sup>

*Bauhinia purpurea* (family: Caesalpiniaceae) is a medium sized, evergreen, ornamental tree found throughout India. The leaves are rigidly sub-coriaceous, glabrous and shallowly cordate. The purple colored flowers of the species distinguishes it specifically from other species of *Bauhinia*. It is presently being used for ailments such as sores, wounds, diarrhea, dropsy, pain, rheumatism, convulsions, delirium, septicemia and so on.<sup>7</sup> Its decoctions are recommended for ulcers as a useful

**\*Corresponding author:**

Email: [shilpapahwa@yahoo.co.in](mailto:shilpapahwa@yahoo.co.in)

wash.<sup>8</sup> The aerial parts of the plant are reported to contain flavanone glycosides, foliar flavonoids, 6-butyl-3-hydroxy flavanone, amino acids, phenyl fatty ester, lutine and  $\beta$ - sitosterol.<sup>9-14</sup> Flavonoids are polyphenolic compounds, widely distributed in the plant kingdom. They are reported to exhibit various pharmacological activities such as CNS activity, cardiogenic activity, lipid-lowering activity, antioxidant activity, hepatoprotective activity, and **hypoglycemic** activity.<sup>15</sup> These active constituents and the above mentioned activities in turn appear to correlate with some other biological activities.

In view of alleged antidiabetic potential of *Bauhinia purpurea*, we have investigated the effect of methanolic extract of its leaf on fasting blood sugar levels in Streptozotocin (STZ) induced diabetic rats. STZ [2-deoxy-2-(3-(methyl-3-nitrosoureido)-D-glucopyranose)] is synthesized by *Streptomyces achromogenes* and is used to induce both insulin dependent and non-insulin dependent DM. STZ is also efficacious after intraperitoneal administration of a dose between 40-60 mg/kg body weight or even a higher dose, but single dose below 40 mg/kg body weight may be ineffective.<sup>16,17</sup>

## MATERIALS AND METHODS

### Plant Material

The leaves of *B. purpurea* were collected from the surrounding areas of Greater Noida in June, 2010. The plant was identified and authenticated (Voucher No NHCP/ NBPGR/2010-20 dated 10th May 2010) by Dr. Anjula Pandey, Principal Scientist, National Bureau of Plant and Genomic Resources (NBPGR), New Delhi, India. A copy of the herbarium had been preserved in the Department of Pharmaceutical Technology, NIET, Greater Noida, India, for future reference. The part was sun dried after washing and then ground to a coarse powder in a mechanical grinder.

### Extraction Procedure

The powdered leaves were macerated with different polar and non polar solvents for seven days, after which each extract was subjected to phytochemical screening using standard methods.<sup>18</sup>

Since the methanolic extract had the maximum number of constituents present in it, methanol was chosen as the solvent for further extraction.

The coarse powder of the leaf (31.83 g) was extracted in a soxhlet apparatus with methanol and

the solvent was removed by controlled evaporation under reduced pressure on a heating mantle at temperature below 60°C. The crude extract thus obtained was tested for its antidiabetic potentiality.

The extract was used to prepare suspensions of 50 mg/kg and 100 mg/kg concentrations using water and Tween 80 : Tween 20 (1:1) for treatment.

### Drugs

STZ was obtained from SRL Lab, Delhi, India. All other chemicals used for the study were of analytical grade.

### Animals

Male Wistar rats weighing between 150 and 200 g were used for antidiabetic activity. Toxicity study was carried out on Albino mice (25-30 g) as per the OECD guidelines and the study was approved by the Institutional Animals Ethics Committee (CPCSEA) (Approval code no. 1121/ac/CPCSEA/07/NIET/IAEC/2010/32P/16). Animals were fed with a standard pellet (Lipton India Ltd., Mumbai, India) and water *ad-libitum*. After randomization into various groups and before initiation of the experiment, the rats were acclimatized for a period of 7 days under 24-28°C temperature, 60-70% relative humidity and 12 h day and night cycle. Animals described as fasted were deprived of food for 16 h but had free access to water.

### Sample Collection

Blood samples were collected by retro-orbital plexus puncture method and blood glucose levels were estimated using an electronic glucometer (Morepen, Baddi, HP, India).

### Effect of *Bauhinia purpurea* Leaf Extract on STZ Induced DM<sup>19</sup>

Diabetes was induced in rats by tail vein injection of STZ (50 mg/kg), dissolved in citrate buffer. One group of 6 identical rats was kept without STZ administration as normal control, group I. Forty eight hours after STZ administration, blood samples were drawn by retro-orbital puncture method and glucose levels were determined to confirm diabetes. The diabetic rats exhibiting blood glucose levels in the range of 140 mg/dl or more were selected for the studies. These diabetic rats were sub divided into 4 groups as follows:

Group II rats, served as diabetic control (STZ induced), were given 0.5 ml of 5% Tween 80 in place of the extract;

Group III diabetic rats were given 50 mg/kg *B. purpurea* leaf extract in 0.5 ml 5% Tween 80;

Group IV diabetic rats were given 100 mg/kg *B. purpurea* leaf extract in 0.5 ml 5% Tween 80;

Group V diabetic rats were given 0.5 ml of 5% Tween 80 containing Glibenclamide (500 µg/kg).

The dose (500 µg/kg) of Glibenclamide was selected based on previous reports.<sup>20</sup> The normal control group of rats (group I) were given 0.5 ml of 5% Tween 80 only.

Each of the control and the test groups consisted of 6 animals. The treatments were continued daily for 15 days. Blood was collected by retro-orbital puncture method for glucose estimation just before drug administration on the 1<sup>st</sup> day and 1 h after drug

flavonoids, phenolic compounds and tannins in the methanolic extract of the part used.

#### Yield

After soxhleting the leaf powder (31.83 g) with methanol, a semisolid, dark, viscous crude extract with 14.54% w/w yield was obtained, which was subjected to further antidiabetic studies.

**The Effect of Methanolic Extract of *Bauhinia purpurea* Leaf on STZ Induced Diabetic Rats**  
Administration of STZ (50 mg/kg, i.p.) led to many folds elevation of fasting blood glucose levels, which was maintained over a period of 2 weeks. Two weeks of daily treatment of methanolic extract of *B. purpurea* led to a dose-dependent fall in blood sugar levels. Vehicle control animals were found to be stable in their body weight but diabetic rats showed significant reduction in body weight during 15 days (**Table 2**). The effect of methanolic leaf extract of *Bauhinia purpurea* on glucose level in STZ induced diabetic rats had been summarized in **Table 3**, where

**Table 1: Preliminary phytochemical screening of various leaf extracts of *Bauhinia purpurea***

S.No	Plant constituents	Extract with solvents						
		Petroleum ether extract	Chloroform extract	Acetic Acid extract	Acetone extract	Methanol extract	Aqueous extract	Benzene extract
1	Alkaloids	-	-	+	-	-	-	+
2	Carbohydrates	-	-	-	-	-	-	-
3	Glycosides	-	+	+	-	+	-	+
4	Saponins	-	-	+	-	+	+	+
5	Phenolic compds & Tanins	+	-	-	+	+	+	-
6	Flavonoids	+	+	-	+	+	+	-
7	Phytosterols	+	+	+	+	+	+	+

administration on days 4, 7 and 10.

#### Statistical Analysis

The results were expressed as Mean ± SEM. Comparison between the groups was made by analysis of variance (ANOVA), followed by Dunnett's test. A value of  $P < 0.001$  was considered significant.<sup>21</sup>

## RESULTS AND DISCUSSION

#### Preliminary Phytochemical Screening

The results of the preliminary phytochemical screening of the various extracts of *B. purpurea* leaf were depicted in **Table 1**, which showed the presence of glycosides, saponins, phytosterols,

a significant reduction ( $P < 0.01$ ) in blood glucose level was observed in the drug treated animals, when compared to the diabetic control group.

The extract has exhibited antidiabetic property in STZ induced diabetic rats, as evident from the glucose levels. The hypoglycemic activity may be ascribed to the presence of flavonoids, which have shown to inhibit cyclooxygenases and promote β-cell regeneration besides having insulin secretory property.<sup>22-24</sup> The results of the present study suggests that the methanolic extract of the leaf of *B. purpurea* illustrates significant hypoglycemic activity, which may be due to the presence of flavonoids in it, as claimed by earlier reports.

**Table 2: Effect of methanolic extract of leaf of *Bauhinia purpurea* (MEBP) on body weight (g) on STZ (50 mg/kg, i.p.) induced diabetes in rats**

Groups	Average body weight (g) of rats				
	1 <sup>st</sup> day	4 <sup>th</sup> day	7 <sup>th</sup> day	10 <sup>th</sup> day	15 <sup>th</sup> day
Normal control (0.5 ml of 5% Tween 80)	182.50 ± 0.53	182.61 ± 0.22	182.67 ± 0.11	183.51 ± 0.45	183.21 ± 0.12
STZ induced diabetic control	200.00 ± 0.12*	182 ± 0.52*	167.54 ± 0.49*	149.22 ± 0.61*	139.11 ± 0.49*
STZ + MEBP (50 mg/kg)	186.61 ± 0.21*	181.57 ± 0.72*	175.00 ± 0.45*	174.40 ± 0.34*	169.62 ± 0.71*
STZ + MEBP (100 mg/kg)	185.53 ± 0.22*	174.13 ± 0.13*	168.67 ± 0.19*	159.31 ± 0.33*	150.00 ± 0.34*
STZ + Glibenclamide (0.5 mg/kg)	189.33 ± 0.65*	186.83 ± 0.63*	181.00 ± 0.55*	178.20 ± 0.22*	176.47 ± 0.57*

Values are mean ± SEM for groups of six animals each

\* P < 0.01 as compared to normal control on corresponding day

**Table 3: Effect of MEBP on glucose level in STZ induced diabetic rats**

Groups	Glucose levels (mg/dl)				
	1 <sup>st</sup> day	4 <sup>th</sup> day	7 <sup>th</sup> day	10 <sup>th</sup> day	15 <sup>th</sup> day
Normal control (0.5 ml of 5% Tween 80)	132.17 ± 0.47	132.17 ± 0.47	131.50 ± 0.42	131.50 ± 0.42	131.50 ± 0.42
STZ induced diabetic control	141.50 ± 0.76*	143.50 ± 1.20**	152.17 ± 0.83*	161.33 ± 0.49*	163.50 ± 0.22*
STZ + MEBP (50 mg/kg)	140.50 ± 0.61*	140.00 ± 0.77**	139.67 ± 0.91*	139.00 ± 1.06*	138.83 ± 1.04*
STZ + MEBP (100 mg/kg)	140.33 ± 0.71*	123.33 ± 16.27**	139.67 ± 0.66*	138.50 ± 1.02*	138.50 ± 1.02*
STZ + Glibenclamide (0.5 mg/kg)	137.83 ± 0.90*	134.33 ± 0.88**	129.00 ± 0.63*	125.17 ± 0.70*	123.67 ± 0.61*

Values are mean ± SEM, 6 rats in each group; STZ (50 mg/kg.) was injected to control and all other drug treated groups \*p<0.01, \*\*p<0.05

**REFERENCES:**

- Vats, V., S.P.Yadav and J.K. Grover, 2004. Ethanolic extract of *Ocimum sanctum* leaves partially attenuates streptozotocin – induced alterations in glycogen content and carbohydrate metabolism in rats. *Journal of Ethnopharmacology*, 1(2): 155-160.
- Kumar, G.P.S., P.Arulselvan, D.S.Kumar and S.P.Subramaniam, 2006. Anti-diabetic activity of fruits of *Terminalia chebula* on Streptozotocin induced diabetic rats. *Journal of health science*, 52(3): 283-291.
- Sumana, G. and S.A. Suryawashi, 2001. Effect of *Vinca rosea* extracts in treatment of alloxan diabetes in male albino rats. *Indian Journal of Experimental biology*, 39: 748-758.
- Holman, R.R. and R.C. Turner, 1991. Oral agents and insulin in the treatment of NIDDM. In *Textbook of Diabetes*, Eds., Pickup, J. and G. Williams. UK : Blackwell, Oxford, pp: 467-469.
- Kameswara, B., M.M.Kesavulu and Ch. Apparao, 2001. Antihyperglycemic activity of *Momordica cymbalaria* in alloxan diabetic rats. *Journal of Ethnopharmacology*, 78 (1): 67-71.
- Ahmed, Mohd.F., Syed Mohd. Kazim, Syed S. Ghori, Syeda S. Mehjabeen, Shaik R. Ahmed, Shaik M.Ali and Mohd. Ibrahim, 2010. Antidiabetic activity of *Vinca rosea* extracts in alloxan-induced diabetic rats. *International Journal of Endocrinology*, 2010:1-6.
- Asolker, L.V., K.K.Kakkar and O.J.Chakre, 2000. 2nd Supplement to Glossary of Indian Medicinal Plants, Part-I (A-K): National Institute of Science Communication, New Delhi.

8. Kirthikar, K.R. and D.B.Basu, 2001. Indian Medicinal Plants: Oriental Enterprises, Dehradun.
9. Salantino, A. and T.T. Blatt, 1999. Foliar flavanoids of nine species of *Bauhinia*. *Revista Brasileira de Botanica*, 22:17-20.
10. Yadava, R.N. and P. Tripathi, 2000. A novel flavones glycoside from the stem of *Bauhinia purpurea*. *Fitotherapia*, 71:88-90.
11. Panda, S and K.Kar, 1999. *Withania somnifera* and *Bauhinia purpurea* in the regulation of circulating thyroid hormone concentrations in female mice. *Journal of Ethnopharmacology*, 67:233-239.
12. Vijayakumari, K.P. and Siddhuraru, 1997. Chemical composition, amino acid content and quality of protein in the legume of *Bauhinia purpurea*. *Journal of Science and Food Agriculture*, 73:279-286.
13. Ramachandran, Reena and B.C.Joshi, 1967. Chemical examination of *Bauhinia purpurea* flowers. *Current Science*, 36:5774-5775.
14. Pettit, G.R., A.Numata, C.Iwamoto, Y. Usami and T.Yamada, 2006. Isolation and structures of bauhiniastatins 1-4 from *Bauhinia purpurea*. *Journal of Natural Products*, 69:323-327.
15. Rajanarayana, K., M.S. Reddy, M.R.Chaluvadi and D.R.Krishna, 2001. Bioflavonoids classification, pharmacological, biochemical effects and therapeutic potential. *Indian Journal of Pharmacology*, 33:2-16.
16. Szkudelski, T, 2001. The mechanism of Alloxan and Streptozotocin action in B cells of the rat pancreas. *Physiol Res*, 50:536-546.
17. Katsumata, K., K.Jr. Katsumata and Y.Katsumata, 1992. Protective effects of diltiazem hydrochloride on the occurrence of alloxan- or streptozotocin induced diabetes in rats. *Hormonal Metabolism*, 24:508-510.
18. Khandelwal, K.R., 2005. Practical Pharmacognosy and Experiments. Nirali Prakashan, Pune.
19. Babu. V., T.Gangadevi and A.Subramonium, 2003. Antidiabetic activity of ethanol extract of *Cassia kleinii* leaf in Streptozotocin –induced diabetic rats and isolation of an active extract and toxicity evaluation of the extract. *Indian Journal of Pharmacology*, 35:290-296.
20. Augusti, K.T. and C.G.Sheela, 1996. Antiperoxide effect of S-allyl cysteine sulfoxide, an insulin secretagogue, in diabetic rats. *Experientia*, 52:115-119.
21. Kulkarni, S.K., 2009. Handbook of Experimental Pharmacology. Vallabh Prakashan, Delhi.
22. Singh, K.N., R.K.Mittal and K.C.Barthwal, 1976. Hypoglycemic activity of *Acacia catechu*, *Acacia suma*, *Albizia odoratissima* seed diets in normal albino rats. *Indian Journal of Medical Research*, 64:754-757.
23. Geeta, B.S., B.C.Mathew and K.C.Augusti, 1994. Hypoglycemic effects of Leucodelphinidin derivative isolated from *Licium bengalenses* (Linn.). *Indian Journal of Physiology and Pharmacology*, 38:220-222.
24. Gupta, S.S., 1994. Prospects and perspectives of natural plant products in medicine. *Indian Journal of Pharmacology*, 26:1-12.